

Onset and duration of 2% lidocaine as inferior alveolar nerve block versus buccal/lingual infiltration of 4% articaine in mandibular second molars: Clinical trial study

Ehsan Esnaashari DDS, MSc¹, Hengameh Bakhtiar DDS, MSc²,
Bahareh Nazari DDS³, Shadi Mirzaei DDS³, Sohrab Tour-Savadkouhi DDS, MSc¹

Original Article

Abstract

BACKGROUND AND AIM: The effectiveness of buccal or lingual (B/L) infiltration of 4% articaine as supplemental injection for pulp anesthesia of mandibular teeth was confirmed in previous studies. However, this study was aimed to compare the effectiveness of 2% lidocaine as inferior alveolar nerve block (IANB) versus B/L infiltration of 4% articaine for pulp anesthesia, as primary injection in mandibular second molars.

METHODS: Thirty adult volunteers ranging from 18 to 40 years old with no systemic disease or medicine intake were included in this split-mouth, double-blind, randomized clinical trial study. Each mandibular side of included subjects was allocated randomly to control group (IANB using 2% lidocaine and 1/80000 epinephrine using direct technique) and B/L infiltration group using 4% articaine (Septanest; Septodont, Saint-Maur-des-Fosses, France). After obtaining base line sensitivity, electric pulp testing (EPT) was done at 5, 8, 11, 15, 20, 25, 30, 45, 60, 75, and 90 minutes post injections. The data were analyzed using chi-square test.

RESULTS: The success rate of anesthesia for IANB group was 83.3% (25 of 30 subjects) and 30% (9 of 30 subjects) for B/L infiltration group, and the difference between the groups was statistically significant ($P = 0.0005$). The mean onset time of pulp anesthesia for IANB group was 22.6 ± 30.9 minutes and 65.5 ± 38.0 for B/L infiltration group, and the difference between the groups was statistically significant ($P = 0.0001$). The mean duration time of pulp anesthesia for IANB group was 53.0 ± 27.4 minutes and 10.6 ± 17.2 for B/L infiltration group, and the difference between the groups was statistically significant ($P = 0.0001$).

CONCLUSION: The results indicated that IANB using 2% lidocaine was more successful than B/L infiltration of 4% articaine in onset and duration of pulp anesthesia of mandibular second molars as primary injections.

KEYWORDS: Articaine; Lidocaine; Local Anesthesia; Molar; Inferior Alveolar Nerve; Volunteers

Citation: Esnaashari E, Bakhtiar H, Nazari B, Mirzaei S, Tour-Savadkouhi S. Onset and duration of 2% lidocaine as inferior alveolar nerve block versus buccal/lingual infiltration of 4% articaine in mandibular second molars: Clinical trial study. *J Oral Health Oral Epidemiol* 2018; 7(3): 113-7.

Predictable anesthesia has an essential role in successful endodontic treatment, and failing to reach this point causes patient miss management. A range of local anesthetic drugs and techniques have been used which lidocaine is the most popular of them, and after a long time articaine was introduced in the United States of America.¹ The inferior

alveolar nerve block (IANB) is the technique of choice for pulpal anesthesia of mandibular teeth. However, the technique is not always successful, and failure rates of 7% to 75% have been reported.²⁻⁴ Therefore, alternatives to this technique were studied in several studies, such as intra ligament, intra osseous, mylohyoid, and infiltration injections.⁵⁻⁸

Articaine is an amide local anesthesia

1- Assistant Professor, Department of Endodontic, School of Dentistry, Tehran Branch, Islamic Azad University, Tehran, Iran

2- Associate Professor, Department of Endodontic, School of Dentistry, Tehran Branch, Islamic Azad University, Tehran, Iran

3- Student of Dentistry, School of Dentistry, Tehran Branch, Islamic Azad University, Tehran, Iran

Correspondence to: Sohrab Tour-Savadkouhi DDS, MSc

Email: s_savadkouhi@yahoo.com

including a thiophene ring, benzene ring, and ester linkage.¹ High lipophilic properties of this solution has made it suitable for infiltration techniques in maxillary and mandibular teeth.⁹⁻¹¹

Several studies have shown the effectiveness of buccal or lingual (B/L) infiltration of 4% articaine as supplemental injection for pulp anesthesia of mandibular teeth.^{9,11} Some studies showed anesthetic effect of B infiltration (BI) of 4% articaine as primary injection on mandibular first molar;^{4,12} however, no one showed its effectiveness as primary injection compared to IANB for pulp anesthesia of second molar.

This study was aimed to compare the effectiveness of 2% lidocaine with 1/80000 epinephrine as IANB versus B/L infiltration of 4% articaine with 1/100000 epinephrine in mandibular second molar pulp anesthesia.

40 years old with no contributing systemic disease or medicine intake, with at least one intact second mandibular molar were included in this split-mouth, double-blind, randomized clinical trial study in Endodontic Department, Dental Branch, Islamic Azad University, Tehran, Iran, from 2015 to 2016 (Figure 1). Exclusion criteria were volunteers younger than 18 or older than 40 years old, allergies to local anesthetics or intolerance of vasoconstrictors, pregnancy, and inability to give informed consent. Sample size was determined based on the results of an initial pilot study on five patients at $\alpha = 0.05$ and a study power of 80%. The Ethics Committee of Islamic Azad University of Tehran approved the study (IR.IAU.DENTAL.REC.1395.19), and informed consent was signed by each volunteer. The proposal of this study was reviewed, confirmed, and recorded in Iranian Registry of Clinical Trials (IRCT) (code No: IRCT2017021523620N7).

Methods

Thirty adult volunteers ranging from 18 to




Figure 1. Consort flow diagram
IANB: Inferior alveolar nerve block; B/L: Buccal/lingual

Each mandibular side of included subjects was randomly (coin flipping method by the patients) allocated to control group (IANB using 2% lidocaine and 1/80000 epinephrine 1.8 ml using direct technique) and the group of B/L infiltration of 4% articaine (Septanest, Septodont, Saint-Maur-des-Fosses, France) using half the solution (0.9 ml) at B and the other half (0.9 ml) at L vestibule just close to the target tooth (second molar). The injections were done in separate sessions by one-week intervals. All the injections were done after negative aspiration at rate of 1 ml/min by one blinded skilled operator; the injections for each volunteer were done by one-week interval.

One of the intact maxillary canines was considered as the control for the pulp tester set up. Base line sensitivity of subjected teeth was determined using electric pulp tester (EPT) (Gentle-Pulse vitality tester; Parkell Inc., Farmingdale, N.Y.) before any injection by blinded trained personnel.

Toothpaste was used as contact media on coronal third of B surface of the crown, and the EPT tip was placed over that. The power was increased incrementally from 1 to 10, until the patient became aware of the electric stimuli. During the EPT test of the teeth, a well-sealed rubber dam was placed over the target tooth (mandibular second molar) to avoid false positive responses due to close contact with neighboring teeth. The pulp testing repeated at 5, 8, 11, 15, 20, 25, 30, 45, 60, 75, and 90 minutes' post injections.⁴

Success in obtaining the pulp anesthesia was considered as at least two subsequent negative responses to EPT in maximum degree. When the pulp anesthesia in subjects was not achieved, the onset was considered

as the maximum follow up time (90 minutes) and the duration as 0.

Comparisons between IANB and B/L infiltration groups for anesthetic success, and onset and duration of pulpal anesthesia were analyzed using chi-square test by SPSS software (version 17, SPSS Inc., Chicago, IL, USA).

Results

All the thirty volunteers (60 subjects which 30 received IANB on one side and 30 received B/L infiltration on the other side) subjected for comparison of success, onset, and duration of pulp anesthesia of mandibular second molars (60% male and 40% female, with mean age of 26.8 ± 1.4).

The success rate of anesthesia for IANB group was 83.3% (25 of 30 subjects) and 30% (9 of 30 subjects) for B/L infiltration group, and the difference between the groups was statistically significant ($P = 0.0005$) (Table 1).

The mean onset time of pulp anesthesia for IANB group was 22.6 ± 30.9 minutes and 65.5 ± 38.0 minutes for B/L infiltration group, and the difference between the groups was statistically significant ($P = 0.0001$) (Figure 2).

The mean duration time of pulp anesthesia for IANB group was 53.0 ± 27.4 minutes and 10.6 ± 17.2 for B/L infiltration group, and the difference between the groups was statistically significant ($P = 0.0001$) (Figure 2).

Discussion

The result of this clinical trial study showed that IANB is advantageous over B/L infiltration of articaine as primary injection in success rate, onset, and duration of pulp anesthesia of mandibular second molars.

Table 1. The success, onset, and duration of pulp anesthesia of mandibular second molar in inferior alveolar nerve block (IANB) and buccal/lingual (B/L) infiltration groups

Groups	Indexes	Success [n (%)]	Failure [n (%)]	Onset (minute) (mean \pm SD)	Duration (minute) (mean \pm SD)
IANB group (n = 30)		25 (83.3)	5 (16.7)	22.6 ± 30.9	53.0 ± 27.4
B/L infiltration group (n = 30)		9 (30.0)	21 (70.0)	65.5 ± 38.0	10.6 ± 17.2
P		0.0005	0.0005	0.0001	0.0001

SD: Standard deviation; IANB: Inferior alveolar nerve block; B/L: Buccal/lingual




Figure 2. Comparing onset and duration of pulp anesthesia of second mandibular molar in inferior alveolar nerve block (IANB) and buccal/lingual (B/L) infiltration groups

Although some studies did not show any significant difference between articaine and lidocaine solutions in obtaining pulp anesthesia of mandibular molars,^{13,14} recent studies found that articaine was remarkably better than lidocaine in pulpal anesthesia of mandibular molars after B/L infiltration.^{11,15,16} For that reason, we analyzed pulpal anesthesia using articaine B/L infiltration for mandibular second molar teeth.

The suspected mechanism of articaine is its better bone-penetration efficacy. Articaine contains a thiophene and benzene ring, which makes the solution to penetrate better through natural barriers.¹⁷ This phenomenon makes the solution suitable for infiltration injections, especially for mandibular molars with thick cortical in both B/L sides.

In a clinical trial study by Jung et al. on mandibular first molar anesthesia after IANB or BI of 4% articaine, they concluded that BI group had faster and more predictable pulp anesthesia at 5 and 8 minutes post injection, and the total success rate was close to IANB group.⁴ The result was totally against our study, and this difference can be explained by the fact that in our study we analyzed pulp anesthesia of second mandibular molars instead of first mandibular molar, which has thicker B cortical bone. Moreover, we used half the articaine solution at B and half at L vestibule, same as Corbett et al.¹² and Foster et al.,⁹ which had used the same technique for obtaining pulp anesthesia of mandibular molars.

In the previous studies, the B/L infiltration of articaine for local anesthesia of mandibular molars was studied as primary or supplementary anesthesia.^{10-12,15,18,19} Most of the studies found B/L infiltration of 4% articaine as an effective technique for pulp anesthesia of mandibular first molars as primary local anesthesia, but the effect was more manifest for B instead of L or B/L injections.¹² However, we found B/L infiltration of articaine as an ineffective technique for pulp anesthesia of mandibular second molars. The success rate of pulp anesthesia of mandibular second molar for this technique was only 30%, which was close to Aggarwal et al. study; however, they had irreversible pulpitis cases, and their pain assessment was during access cavity preparation and root canal instrumentation instead of normal pulp cases and EPT sensibility test in this study.²⁰ The higher success rate of IANB using 2% lidocaine in this study compared to other studies can be explained by using normal healthy dental pulps instead of teeth with irreversible pulpitis subjected to other studies.^{19,21}

One of the main concerns, when using articaine as local anesthesia especially in mandibular nerve block, is the paresthesia as an important side effect.²² In a comprehensive review study on articaine as local anesthesia by Kakroudi et al., it was concluded that the solution appears safe and the adverse effects are very rare.²³ One of the main limitations of this study was non-inflamed and normal pulp of subjects for maximum unification, while most of the anesthetic failures suspected in inflamed pulp conditions.

Conclusion

It seems that IANB using 2% lidocaine was more effective than B/L infiltration of 4% articaine in success, onset, and duration of pulp anesthesia of mandibular second molars as primary injections.

Conflict of Interests

Authors have no conflict of interest.

Acknowledgments

The authors acknowledge generous support

by research committee of Dental school, Islamic Azad University, Tehran, Iran.

References

1. Malamed SF, Gagnon S, Leblanc D. Articaine hydrochloride: A study of the safety of a new amide local anesthetic. *J Am Dent Assoc* 2001; 132(2): 177-85.
2. Goodman A, Reader A, Nusstein J, Beck M, Weaver J. Anesthetic efficacy of lidocaine/meperidine for inferior alveolar nerve blocks. *Anesth Prog* 2006; 53(4): 131-9.
3. Kanaa MD, Meechan JG, Corbett IP, Whitworth JM. Speed of injection influences efficacy of inferior alveolar nerve blocks: A double-blind randomized controlled trial in volunteers. *J Endod* 2006; 32(10): 919-23.
4. Jung IY, Kim JH, Kim ES, Lee CY, Lee SJ. An evaluation of buccal infiltrations and inferior alveolar nerve blocks in pulpal anesthesia for mandibular first molars. *J Endod* 2008; 34(1): 11-3.
5. Nusstein J, Claffey E, Reader A, Beck M, Weaver J. Anesthetic effectiveness of the supplemental intraligamentary injection, administered with a computer-controlled local anesthetic delivery system, in patients with irreversible pulpitis. *J Endod* 2005; 31(5): 354-8.
6. Bigby J, Reader A, Nusstein J, Beck M, Weaver J. Articaine for supplemental intraosseous anesthesia in patients with irreversible pulpitis. *J Endod* 2006; 32(11): 1044-7.
7. Stein P, Brueckner J, Milliner M. Sensory innervation of mandibular teeth by the nerve to the mylohyoid: Implications in local anesthesia. *Clin Anat* 2007; 20(6): 591-5.
8. Srinivasan N, Kavitha M, Loganathan CS, Padmini G. Comparison of anesthetic efficacy of 4% articaine and 2% lidocaine for maxillary buccal infiltration in patients with irreversible pulpitis. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2009; 107(1): 133-6.
9. Foster W, Drum M, Reader A, Beck M. Anesthetic efficacy of buccal and lingual infiltrations of lidocaine following an inferior alveolar nerve block in mandibular posterior teeth. *Anesth Prog* 2007; 54(4): 163-9.
10. McEntire M, Nusstein J, Drum M, Reader A, Beck M. Anesthetic efficacy of 4% Articaine with 1:100,000 epinephrine versus 4% articaine with 1:200,000 epinephrine as a primary buccal infiltration in the mandibular first molar. *J Endod* 2011; 37(4): 450-4.
11. Robertson D, Nusstein J, Reader A, Beck M, McCartney M. The anesthetic efficacy of articaine in buccal infiltration of mandibular posterior teeth. *J Am Dent Assoc* 2007; 138(8): 1104-12.
12. Corbett IP, Kanaa MD, Whitworth JM, Meechan JG. Articaine infiltration for anesthesia of mandibular first molars. *J Endod* 2008; 34(5): 514-8.
13. Aggarwal V, Singla M, Miglani S. Comparative evaluation of anesthetic efficacy of 2% lidocaine, 4% articaine, and 0.5% bupivacaine on inferior alveolar nerve block in patients with symptomatic irreversible pulpitis: A prospective, randomized, double-blind clinical trial. *J Oral Facial Pain Headache* 2017; 31(2): 124-8.
14. Corbella S, Taschieri S, Mannocci F, Rosen E, Tsesis I, Del Fabbro M. Inferior alveolar nerve block for the treatment of teeth presenting with irreversible pulpitis: A systematic review of the literature and meta-analysis. *Quintessence Int* 2017; 48(1): 69-82.
15. Martin M, Nusstein J, Drum M, Reader A, Beck M. Anesthetic efficacy of 1.8 mL versus 3.6 mL of 4% articaine with 1:100,000 epinephrine as a primary buccal infiltration of the mandibular first molar. *J Endod* 2011; 37(5): 588-92.
16. da Silva-Junior GP, de Almeida Souza LM, Groppo FC. Comparison of Articaine and lidocaine for buccal infiltration after inferior alveolar nerve block for intraoperative pain control during impacted mandibular third molar surgery. *Anesth Prog* 2017; 64(2): 80-4.
17. Zain M, Rehman Khattak SU, Sikandar H, Shah SA, Fayyaz. Comparison of anaesthetic efficacy of 4% articaine primary buccal infiltration versus 2% lidocaine inferior alveolar nerve block in symptomatic mandibular first molar teeth. *J Coll Physicians Surg Pak* 2016; 26(1): 4-8.
18. Kanaa MD, Whitworth JM, Corbett IP, Meechan JG. Articaine and lidocaine mandibular buccal infiltration anesthesia: A prospective randomized double-blind cross-over study. *J Endod* 2006; 32(4): 296-8.
19. Matthews R, Drum M, Reader A, Nusstein J, Beck M. Articaine for supplemental buccal mandibular infiltration anesthesia in patients with irreversible pulpitis when the inferior alveolar nerve block fails. *J Endod* 2009; 35(3): 343-6.
20. Aggarwal V, Singla M, Kabi D. Comparative evaluation of anesthetic efficacy of Gow-Gates mandibular conduction anesthesia, Vazirani-Akinosi technique, buccal-plus-lingual infiltrations, and conventional inferior alveolar nerve anesthesia in patients with irreversible pulpitis. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2010; 109(2): 303-8.
21. Parirokh M, Satvati SA, Sharifi R, Rekabi AR, Gorjestani H, Nakhaee N, et al. Efficacy of combining a buccal infiltration with an inferior alveolar nerve block for mandibular molars with irreversible pulpitis. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2010; 109(3): 468-73.
22. Piccinni C, Gissi DB, Gabusi A, Montebugnoli L, Poluzzi E. Paraesthesia after local anaesthetics: An analysis of reports to the FDA Adverse Event Reporting System. *Basic Clin Pharmacol Toxicol* 2015; 117(1): 52-6.
23. Kakroudi SH, Mehta S, Millar BJ. Articaine hydrochloride: Is it the solution? *Dent Update* 2015; 42(1): 88-3.