

## The effect of periodontal status on the TSH and FT4 levels

Devrim Deniz Üner , Bozan Serhat Izol


DOI: <https://doi.org/10.34172/johoe.2202.1456>

### Article History:

Received Date: February 19, 2022

Accepted Date: June 28, 2023

published Author Accepted Version: May 8, 2024



**Copyright:** © 2024 The Author(s); Published by Kerman University of Medical Sciences. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Please cite this article as:** Alwan A, Jamison DT, Siddiqi S, Vassall A. Pakistan's progress on universal health coverage: lessons learned in priority setting and challenges ahead in reinforcing primary healthcare. *Int J Health Policy Manag.* 2024;x(x):x-x. doi.org/10.34172/johoe.2202.1456

**This PDF file is an Author Accepted Manuscript (AAM) version, which has not been typeset or copyedited, but has been peer reviewed. JOHOE publishes the AAM version of all accepted manuscripts upon acceptance to reach fast visibility. During the proofing process, errors may be discovered (by the author/s or editorial office) that could affect the content, and we will correct those in the final proof.**



Manuscript Type: Original Article

## **The effect of periodontal status on the TSH and FT4 levels**

Devrim Deniz Üner\*<sup>1</sup>, Bozan Serhat Izol<sup>2</sup>

1. Department of Periodontology, Faculty of Dentistry, Harran University, Şanlıurfa, Turkey.
2. Department of Periodontology, Faculty of Dentistry, Bingöl University, Bingöl, Turkey.

**Corresponding Author:** Dr. Devrim Deniz Üner; [dvrmdnznr@gmail.com](mailto:dvrmdnznr@gmail.com)

### **Abstract**

**Background:** Some bacteria and viruses cause thyroid gland inflammation. Periodontal diseases lead to the production of numerous bacteria in the mouth that spread to other tissues and organs, causing infection in those sites. This investigation's goal was to the correlation between thyroid abnormalities with periodontal disorders.

**Methods:** The research study was carried out on 96 patients who provided blood samples from the total of 1012 applicants who enrolled to the periodontology clinic. The patients were categorized into 4 categories based on their age: (i) ages ranging 0-20, (ii) ages ranging 21-40, (iii) ages ranging 41-60, and (iv) ages over 61. In addition, they were additionally divided into three groups according to both TSH and FT4 levels: The TSHa and FT4a group included patients with below-normal values, the TSHb and FT4b group included patients with normal TSH and FT4 values, and the TSHc and FT4c group included patients with above-normal TSH and FT4 values. The statistical analysis was performed using SPSS 21. Frequency analysis and the Mann-Whitney U Analyses were utilized for statistical analysis., and the level of significance was set at  $P < 0.05$ .

**Results:** The median TSH and FT4 levels of the patients were  $1.98 \pm 1.28$  mIU/L and  $1.19 \pm 0.32$  mIU/L, in that order there were no significant statistical differences seen in TSH and FT4 levels between genders or age groups. According to their FT4 and TSH levels, 7.3% of the patients had subclinical hypothyroidism and 4.2% had subclinical hyperthyroidism. In total, 11.5% of the patients had thyroid dysfunction.

**Conclusion:** The results indicated that thyroid dysfunction was present in a remarkably high proportion of patients (11.5%).

**Keywords:** thyroid dysfunction, gingivitis, periodontitis, periodontal health

### **Introduction**

The thyroid gland a distinctive butterfly form, consisting of two lobes on the both of two sides, which are merged in the center by a thin structure known as the 'isthmus'.<sup>1, 2</sup> The

gland is located corresponding to the vertebral levels C5-T1 and is wrapped around the trachea anteriorly.<sup>2</sup> In reaction to thyroid stimulating hormone (TSH), the thyroid gland releases free thyroxine hormone (FT4). The secreted FT4 is converted to triiodothyronine (FT3), which is a stronger hormone, by deiodinase enzymes.<sup>3</sup> Hyper- or hypo-thyroidism is just one of many physical and mental health issues that can develop from various thyroid diseases, which also include benign, functional, inflammatory, autoimmune, and neoplastic disorders.<sup>4-7</sup> TSH and FT4 are highly sensitive indicators of thyroid function and management of hypo- and hyper-thyroidism.<sup>8</sup> In healthy individuals, normal TSH values vary between 0.4 and 4.0 mIU/ml<sup>9</sup> and normal FT4 values vary between 0.9 and 1.592 ng/dL.<sup>10</sup> Abnormal TSH and FT4 levels result in hyperthyroidism, subclinical hyperthyroidism, hypothyroidism, and subclinical hypothyroidism, which are generally referred to as thyroid dysfunction.<sup>11</sup>

Periodontitis is an infectious condition that is defined by specific characteristics by the spread of inflammation.<sup>12</sup> The correlation between periodontal disorders and numerous systemic issues have been documented in the literature.<sup>13, 14</sup>

Acute suppurative thyroiditis (AST), an infection in the thyroid gland, is a rare infectious disease characterized by high FT4 and low TSH values, primarily responsible by bacteria for example *S. aureus*, *Strep. pyogenes*, *S. epidermidis*, and *Strep. pneumoniae*.<sup>15</sup> *Strep. anginosus*, a type of bacteria causing head and neck abscesses, is frequently seen in the oral region and also causes AST.<sup>15, 16</sup> In a 2013 study, Kumagai et al. detected *Streptococcus anginosus* in 28 (75.6%) out of 37 periodontitis patients and only in three (15%) out of 20 healthy patients.<sup>17</sup> The present investigation was conducted to examine in the event TSH and FT4 levels of patients with oral infection.

## Methods

**Patient selection** :This study evaluated the blood test results of patients who provided the results of a blood test up to 30 days prior to the diagnosis of gingivitis or periodontitis in Harran University Faculty of Dentistry Periodontology Clinic between 2017 and 2020. The patients were selected without regard to the severity of periodontitis or gingivitis. Among the patients whose blood tests were accessed via hospital databases, the patients whose TSH and FT4 values were quantified in their blood tests were utilized in research. The study participants were categorized into four age groups: (i) ages ranging 0-20, (ii) ages ranging 21-40, (iii) ages ranging 41-60, and (iv) ages over 61.

The patients were assessed in three distinct groups: The TSHa and FT4a group included patients with below-normal values, the TSHb and FT4b group included patients with normal TSH and FT4 values, and the TSHc and FT4c group included patients with above-

normal TSH and FT4 values. The study was approved by the Harran University Clinical Research Ethics Committee on March 1, 2021, with the reference number 15271.

### **Statistical analysis**

The data were examined utilizing SPSS statistical software for PCs edition 22.0, developed by IBM Corp. and located in Armonk, NY. The data underwent evaluation for normal distribution via the Kolmogorov-Smirnov test, while the homogeneity of variance was assessed through Levene's test. The Mann-Whitney U test was used to compare variables that have a distribution that is not normal. A P-value below 0.05 was deemed statistically significant.

### **Results**

Out of the total of 1012 patients who applied to our periodontology clinic, 96 provided blood test results for the assessment of thyroid function. All the 96 patients had a TSH value in the hospital databases, while only 84 of them had a FT4 value.

The 96 patients whose TSH values were available in the hospital databases comprised 50 males and 46 females with an average age of  $42.74 \pm 16.27$  for males and  $38.07 \pm 12.79$  for females. In these patients, the median TSH value was 1.55 mIU/L with a range, 0.35–4.48 mIU/L for men and 1.90 mIU/L with a range, 0.23–6.88 mIU/L for women, and there is no discernible disparity was observed among the genders (Figure 1). By contrast, median FT4 value was 1.17 mIU/L with a range, 0.77–3.67 mIU/L for female and 1.11 mIU/L with a range, 0.83–1.77 mIU/L for female, and there is no discernible disparity was noticed in the context of the genders (Figure 2). In the analysis of TSH values with regard to age groups, the median TSH value was 1.55 mIU/L with a range, 1.27–4.48 mIU/L in group I, 1.78 mIU/L with a range, 0.23–6.86 mIU/L in group II, 1.93 mIU/L with a range, 0.27–6.88 mIU/L in group III, and 1.17 mIU/L with a range, 0.28–3.44 mIU/L in group IV. As for FT4 values, the median FT4 value was 1.18 mIU/L with a range, 1.10–1.27 mIU/L in group I, 1.14 mIU/L with a range, 0.83–3.67 mIU/L in group II, 1.17 mIU/L with a range, 0.80–1.77 mIU/L in group III, and 1.07 mIU/L with a range, 0.77–1.20 mIU/L in group IV (Table 1). As a result of the analysis, the P value was calculated as 0.401 for TSH and 0.69 for FT4. The analyses confirmed that there was not a noticeable variation in the TSH and FT4 levels. (Figure 3). Furthermore, in all patients, the mean TSH value was  $1.98 \pm 1.28$  mIU/L and the mean FT4 value was  $1.19 \pm 0.32$  mIU/L. Based on their TSH values, 4 (4.2%) out of 96 patients were classified into the TSHb group, 85 (88.5%) were assigned to the TSHb group, and 7 (7.3%) were assigned to the TSHc group (Table 2). In contrast, based on their FT4 values, 7 (8.3%) patients were assigned to the FT4a group, 76 (90.5%) were

assigned to the FT4b group, and 1 (1.12%) patient were assigned to the FT4c group (Table 3).

## Discussion

The present study evaluated individuals who applied to our periodontology clinic and were diagnosed with gingivitis or periodontitis and whose serum TSH and FT4 values that had been obtained within the last one month were available in hospital databases. In the analysis, the median TSH and FT4 levels were  $1.98 \pm 1.28$  mIU/L and  $1.19 \pm 0.32$  mIU/L, respectively. Additionally, there was no discernible disparity observed between genders and among age groups with regard to TSH and FT4 levels. According to their FT4 and TSH levels, 7.3% of the patients were subclinical hypothyroidism, 4.2% of them had subclinical hyperthyroidism, and 11.5% of them had thyroid dysfunction.

Assessing the TSH level is the optimal method for evaluating thyroid function. Approximately 95% of the healthy population have a serum TSH value ranging from 0.4 to 2.5 mU/L.<sup>18, 19</sup> The thyroid gland becomes dysfunctional along with aging, particularly in women. Additionally, several studies indicated that the amounts of TSH in the blood rise with aging.<sup>20, 21</sup> Of note, Lee et al. found no significant difference among individuals aged below 60 years with regard to serum TSH values.<sup>22</sup> Conversely, several research have demonstrated that gender, age, and race exert an influence on serum TSH levels.<sup>23-25</sup> There was no statistically significant variation observed in serum TSH and FT4 levels across age categories in our research.

The relationship among serum TSH and FT4 levels and systemic diseases and other medical disorders has been the subject of several researches.<sup>14, 22, 26-30</sup> Albrecht et al. evaluated the association among focus lack hyperactivity condition and FT3 in 2020 and reported a positive relationship.<sup>31</sup> In addition, some other studies reported BMI (body mass index) is positively correlated with TSH levels and also noted that TSH levels were significantly higher in obese kids compared to kids that are of a healthy weight.<sup>32, 33</sup> On the other hand, given that thyroid hormones act a crucial function on the neurological progress during the fetal period, several research have examined the correlation between psychiatric condition and thyroid hormones and reported that patients with hyperthyroidism had a higher hospitalization rate for psychiatric treatment compared to healthy individuals.<sup>6</sup> Moreover, Bauer et al. found that hip and vertebral fractures were observed more frequently in women aged over 65 years with low levels of TSH in the blood.<sup>34</sup>

Thyroiditis can cause homeostatic imbalance and affect the healing capacity of tissues.<sup>35</sup>,<sup>36</sup> Nevertheless, the majority of these research have evaluated the association among thyroid disorders with oral lichen planus, and only a minority of them have investigated the correlation between periodontal status and thyroid diseases.<sup>14, 37-40</sup> In a 2017 review,

Li et al. demonstrated the incidence of thyroid disorders was greater among individuals having oral lichen planus.<sup>37</sup> In another review, Aldulaijan et al. reported on a relationship between periodontitis and hypothyroidism.<sup>14</sup> Additionally, there have been reports that uncontrolled thyroid diseases result in periodontal destruction.<sup>41</sup> Another study suggested that non-surgical periodontal treatment affects serum levels of thyroid hormones.<sup>42</sup> Conversely, Tuysuz and Beker stated that the occurrence of subclinical hypothyroidism in grown-ups with Down syndrome was 25.3%.<sup>43</sup> In some other studies, hyperthyroidism was detected in approximately 4.4% among those diagnosed with type 2 DM, while subclinical hyperthyroidism was detected in 2–4% of the individuals.<sup>44</sup> Conversely, a separate study found that the occurrence of thyroid malfunction in patients have Type 2 diabetes was 16.2%.<sup>45</sup>

The patient ratio in our research who provided a blood specimen for the assessment of serum TSH and FT4 values one month before gingival examination was 9.48% among the total patients who were examined. The median TSH value was  $1.98 \pm 1.28$  mIU/L and the median FT4 value was  $1.19 \pm 0.32$  mIU/L. Additionally, when the TSH results of the subjects in the research were analyzed, it was concluded that four (4.2%) patients whose TSH levels were below the usual range had normal FT4 levels and these individuals were categorized as having subclinical hypothyroidism. However, it was observed that seven (7.3%) patients whose TSH levels exceeded the upper bound had normal FT4 levels. Overall, these findings indicate that thyroid dysfunction was observed in 11.5% ( $4.2\% \pm 7.3\%$ ) of our patients.

### **Strengths and Limitations**

Although this study explains the relationship between thyroid diseases and periodontal diseases, there are certain constraints associated with it. The primary constraint is that the change in the systemic status of the patients whose blood tests were obtained in the last month is not known exactly.

### **Conclusion**

The results indicated a remarkably high prevalence of thyroid dysfunction (11.5%) in periodontal problems.

### **Acknowledgments**

We are thankful for your support to the editors and employees of Journal of Oral Health and Oral Epidemiology for their support of our study.

### **Authors' Contribution**

**Conceptualization:** D.D. Üner, B.S. İzol  
**Data Curation:** D.D. Üner, B.S. İzol  
**Investigation:** D.D. Üner, B.S. İzol  
**Formal Analysis:** D.D. Üner, B.S. İzol  
**Methodology:** D.D. Üner  
**Project Administration:** D.D. Üner, B.S. İzol  
**Supervision:** D.D. üner  
**Software:** Devrim Deniz Üner  
**Resource:** D.D. Üner, B.S. İzol  
Validation: D.D. Üner, B.S. İzol  
**Visualization:** D.D. Üner, B.S. İzol  
**Writing- Original Draft:** D.D. Üner  
**Writing- Review & Editing:** D.D. Üner

### Competing Interests

The writers of this work have clearly stated that they have no conflicts of interest.

### Data Availability Statement

Nil.

### Ethical Approval

The Harran University Clinical Research Ethics Committee granted clearance for this study with the number 15271 on March 1, 2021.

### Funding

There was no explicit support or funding from any governmental, corporate, or philanthropic organizations for this research.

### References

1. Hoda SA, Hoda RS. Robbins and Cotran Pathologic Basis of Disease. American Journal of Clinical Pathology. 2020;154(6):869-. doi: 10.1093/ajcp/aqaa163
2. Morphologia E. Junqueira's Basic Histology : Text & Atlas (15th ed.), 2018. Morphologia. 2019;13:101-4. doi: 10.26641/1997-9665.2019.3.101-104
3. Leung AM, Braverman LE. Consequences of excess iodine. Nature Reviews Endocrinology. 2014;10(3):136-42. doi: 10.1038/nrendo.2013.251
4. Ragusa F, Fallahi P, Elia G, Gonnella D, Paparo SR, Giusti C, et al. Hashimotos' thyroiditis: Epidemiology, pathogenesis, clinic and therapy. Best Practice & Research Clinical Endocrinology & Metabolism. 2019;33(6):101367. doi: 10.1016/j.beem.2019.101367

5. Muller I, Moran C, Lecumberri B, Decallonne B, Robertson N, Jones J, Dayan CM. 2019 European Thyroid Association guidelines on the management of thyroid dysfunction following immune reconstitution therapy. *European thyroid journal*. 2019;8(4):173-85. doi: 10.1159/000500881
6. Brandt F, Thvilum M, Almind D, Christensen K, Green A, Hegedüs L, Brix TH. Hyperthyroidism and psychiatric morbidity: evidence from a Danish nationwide register study. *European Journal of Endocrinology*. 2014;170(2):341-8. doi: 10.1530/EJE-13-0708
7. Thvilum M, Brandt F, Almind D, Christensen K, Brix TH, Hegedüs L. Type and extent of somatic morbidity before and after the diagnosis of hypothyroidism. A nationwide register study. *PLoS One*. 2013;8(9):e75789. doi: 10.1371/journal.pone.0075789
8. Liyanage Y, Siriwardhana I, Dissanayake M, Dayanath B. Study on diurnal variation in TSH and freeT4 levels of healthy adults. *Sri Lanka Journal of Diabetes Endocrinology and Metabolism*. 2018;8(1). doi: 10.4038/sjdem.v8i1.7346
9. Paragliola RM, Di Donna V, Locantore P, Papi G, Pontecorvi A, Corsello SM. Factors Predicting Time to TSH Normalization and Persistence of TSH Suppression After Total Thyroidectomy for Graves' Disease. *Front Endocrinol (Lausanne)*. 2019;10:95. doi: 10.3389/fendo.2019.00095
10. Murillo-Llorente MT, Fajardo-Montañana C, Pérez-Bermejo M. Reference Values of Thyroid Hormones During the First Trimester of Pregnancy in Valencian Community (Spain) and Their Relationship with Iodine Intake. *Nutrients*. 2020;12(5). doi: 10.3390/nu12051433
11. Fatourechi V, Klee GG, Grebe SK, Bahn RS, Brennan MD, Hay ID, et al. Effects of reducing the upper limit of normal TSH values. *Jama*. 2003;290(24):3195-6. doi: 10.1001/jama.290.24.3195-b
12. Tonetti MS, Greenwell H, Kornman KS. Staging and grading of periodontitis: Framework and proposal of a new classification and case definition. *J Periodontol*. 2018;89 Suppl 1:S159-s72. doi: 10.1002/jper.18-0006
13. Genco RJ, Borgnakke WS. Risk factors for periodontal disease. *Periodontol 2000*. 2013;62(1):59-94. doi: 10.1111/j.1600-0757.2012.00457.x
14. Aldulaijan HA, Cohen RE, Stellrecht EM, Levine MJ, Yerke LM. Relationship between hypothyroidism and periodontitis: A scoping review. *Clin Exp Dent Res*. 2020;6(1):147-57. doi: 10.1002/cre2.247



15. Noguchi S, Yatera K, Kawanami T, Yamasaki K, Naito K, Akata K, et al. The clinical features of respiratory infections caused by the Streptococcus anginosus group. *BMC Pulm Med*. 2015;15:133. doi: 10.1186/s12890-015-0128-6
16. Wu C, Zhang Y, Gong Y, Hou Y, Li S, Zou Y, Ge J. Two cases of bacterial suppurative thyroiditis caused by Streptococcus anginosus. *Endocr Pathol*. 2013;24(1):49-53. doi: 10.1007/s12022-013-9236-0
17. Kumagai K, Sugano N, Takane M, Iwasaki H, Tanaka H, Yoshinuma N, et al. Detection of Streptococcus anginosus from saliva by real-time polymerase chain reaction. *Lett Appl Microbiol*. 2003;37(5):370-3. doi: 10.1046/j.1472-765x.2003.01405.x
18. Baloch Z, Carayon P, Conte-Devolx B, Demers LM, Feldt-Rasmussen U, Henry JF, et al. Laboratory medicine practice guidelines. Laboratory support for the diagnosis and monitoring of thyroid disease. *Thyroid*. 2003;13(1):3-126. doi: 10.1089/105072503321086962
19. Pearce SH, Brabant G, Duntas LH, Monzani F, Peeters RP, Razvi S, Wemeau JL. 2013 ETA Guideline: Management of Subclinical Hypothyroidism. *Eur Thyroid J*. 2013;2(4):215-28. doi: 10.1159/000356507
20. Surks MI, Hollowell JG. Age-specific distribution of serum thyrotropin and antithyroid antibodies in the US population: implications for the prevalence of subclinical hypothyroidism. *J Clin Endocrinol Metab*. 2007;92(12):4575-82. doi: 10.1210/jc.2007-1499
21. Bremner AP, Feddema P, Leedman PJ, Brown SJ, Beilby JP, Lim EM, et al. Age-related changes in thyroid function: a longitudinal study of a community-based cohort. *J Clin Endocrinol Metab*. 2012;97(5):1554-62. doi: 10.1210/jc.2011-3020
22. Lee YK, Kim JE, Oh HJ, Park KS, Kim SK, Park SW, et al. Serum TSH level in healthy Koreans and the association of TSH with serum lipid concentration and metabolic syndrome. *Korean J Intern Med*. 2011;26(4):432-9. doi: 10.3904/kjim.2011.26.4.432
23. Boucai L, Surks MI. Reference limits of serum TSH and free T4 are significantly influenced by race and age in an urban outpatient medical practice. *Clin Endocrinol (Oxf)*. 2009;70(5):788-93. doi: 10.1111/j.1365-2265.2008.03390.x
24. Beydoun MA, Beydoun HA, Kitner-Triolo MH, Kaufman JS, Evans MK, Zonderman AB. Thyroid hormones are associated with cognitive function: moderation by sex, race, and depressive symptoms. *J Clin Endocrinol Metab*. 2013;98(8):3470-81. doi: 10.1210/jc.2013-1813

25. Surks MI, Boucai L. Age- and race-based serum thyrotropin reference limits. *J Clin Endocrinol Metab.* 2010;95(2):496-502. doi: 10.1210/jc.2009-1845
26. Svare A, Nilsen TI, Bjoro T, Asvold BO, Langhammer A. Serum TSH related to measures of body mass: longitudinal data from the HUNT Study, Norway. *Clin Endocrinol (Oxf).* 2011;74(6):769-75. doi: 10.1111/j.1365-2265.2011.04009.x
27. Chen C, Xia F, Chen Y, Zhang K, Cheng J, Li Q, et al. Association Between Thyroid-Stimulating Hormone and Renal Function: a Mendelian Randomization Study. *Kidney Blood Press Res.* 2018;43(4):1121-30. doi: 10.1159/000491808
28. Kats L, Goldman Y, Kahn A, Goldman V, Gorsky M. Oral lichen planus and thyroid gland diseases: possible associations. *BMC Oral Health.* 2019;19(1):169. doi: 10.1186/s12903-019-0859-5
29. Jayanthi R, Srinivasan AR, Hanifah M, Maran AL. Associations among Insulin Resistance, Triacylglycerol/High Density Lipoprotein (TAG/HDL ratio) and Thyroid hormone levels-A study on Type 2 diabetes mellitus in obese and overweight subjects. *Diabetes Metab Syndr.* 2017;11 Suppl 1:S121-s6. doi: 10.1016/j.dsx.2016.12.020
30. Şandru F, Carsote M, Petca RC, Gheorghisan-Galateanu AA, Petca A, Valea A, Dumitraşcu MC. COVID-19-related thyroid conditions (Review). *Exp Ther Med.* 2021;22(1):756. doi: 10.3892/etm.2021.10188
31. Albrecht D, Ittermann T, Thamm M, Grabe HJ, Bahls M, Völzke H. The association between thyroid function biomarkers and attention deficit hyperactivity disorder. *Sci Rep.* 2020;10(1):18285. doi: 10.1038/s41598-020-75228-w
32. Ünüvar T, Anık A, Catlı G, Esen I, Abacı A, Büyükgebiz A, Böber E. Isolated hyperthyrotropinemia in childhood obesity and its relation with metabolic parameters. *J Endocrinol Invest.* 2014;37(9):799-804. doi: 10.1007/s40618-014-0100-y
33. Wolters B, Lass N, Reinehr T. TSH and free triiodothyronine concentrations are associated with weight loss in a lifestyle intervention and weight regain afterwards in obese children. *Eur J Endocrinol.* 2013;168(3):323-9. doi: 10.1530/eje-12-0981
34. Bauer DC, Ettinger B, Nevitt MC, Stone KL. Risk for fracture in women with low serum levels of thyroid-stimulating hormone. *Ann Intern Med.* 2001;134(7):561-8. doi: 10.7326/0003-4819-134-7-200104030-00009
35. Venkatesh Babu NS, Patel PB. Oral health status of children suffering from thyroid disorders. *J Indian Soc Pedod Prev Dent.* 2016;34(2):139-44. doi: 10.4103/0970-4388.180443

36. McAninch EA, Bianco AC. Thyroid hormone signaling in energy homeostasis and energy metabolism. *Ann N Y Acad Sci.* 2014;1311:77-87. doi: 10.1111/nyas.12374
37. Li D, Li J, Li C, Chen Q, Hua H. The Association of Thyroid Disease and Oral Lichen Planus: A Literature Review and Meta-analysis. *Front Endocrinol (Lausanne).* 2017;8:310. doi: 10.3389/fendo.2017.00310
38. Guarneri F, Giuffrida R, Di Bari F, Cannavò SP, Benvenga S. Thyroid Autoimmunity and Lichen. *Front Endocrinol (Lausanne).* 2017;8:146. doi: 10.3389/fendo.2017.00146
39. Tang Y, Shi L, Jiang B, Zhou Z, Shen X. A Cross-Sectional Study of Oral Lichen Planus Associated With Thyroid Diseases in East China. *Front Endocrinol (Lausanne).* 2019;10:928. doi: 10.3389/fendo.2019.00928
40. Robledo-Sierra J, Landin-Wilhelmsen K, Nyström HF, Mattsson U, Jontell M. Clinical characteristics of patients with concomitant oral lichen planus and thyroid disease. *Oral Surg Oral Med Oral Pathol Oral Radiol.* 2015;120(5):602-8. doi: 10.1016/j.oooo.2015.08.001
41. Zahid TM, Wang BY, Cohen RE. The effects of thyroid hormone abnormalities on periodontal disease status. *J Int Acad Periodontol.* 2011;13(3):80-5. doi:
42. Bhanekar RR, Hungund S, Kambalyal P, Singh V, Jain K. Effect of nonsurgical periodontal therapy on thyroid stimulating hormone in hypothyroid patients with periodontal diseases. *Indian J Dent Res.* 2017;28(1):16-21. doi: 10.4103/ijdr.IJDR\_174\_16
43. Tüysüz B, Beker DB. Thyroid dysfunction in children with Down's syndrome. *Acta Paediatr.* 2001;90(12):1389-93. doi: 10.1080/08035250152708770
44. Biondi B, Kahaly GJ, Robertson RP. Thyroid Dysfunction and Diabetes Mellitus: Two Closely Associated Disorders. *Endocr Rev.* 2019;40(3):789-824. doi: 10.1210/er.2018-00163
45. Jali MV, Kambar S, Jali SM, Pawar N, Nalawade P. Prevalence of thyroid dysfunction among type 2 diabetes mellitus patients. *Diabetes Metab Syndr.* 2017;11 Suppl 1:S105-s8. doi: 10.1016/j.dsx.2016.12.017

## Tables

**Table 1.** Age-dependent distribution of TSH and FT4 concentrations

Group		TSH	FT4
1	N	8	8
	Median	1.55	1.18
	Minimum	1.27	1.10
	Maximum	4.48	1.27
2	N	39	32
	Median	1.78	1.14
	Minimum	0.23	0.83
	Maximum	6.86	3.67
3	N	39	36
	Median	1.93	1.17
	Minimum	0.27	0.80
	Maximum	6.88	1.77
4	N	10	8
	Median	1.17	1.07
	Minimum	0.28	0.77
	Maximum	3.44	1.20
Total	N	96	84
	Median	1.76	1.15
	Minimum	0.23	0.77
	Maximum	6.88	3.67

**Table 2.** Patients' distribution in accordance with TSH categories.


	Frequency	%
TSHa	4	4.2
TSHb	85	88.5
TSHc	7	7.3
Total	96	100

**Table 3.** Patients' distribution in accordance with FT4 categories.


	Frequency	%
FT4a	7	8.3
FT4b	76	90.5
FT4c	1	1.2
Total	84	100

**Figures**

**Figure 1.** TSH levels according to genders



**Figure 2.** Distribution of FT4 levels according to genders



**Figure 3.** Average values of TSH-FT4 levels 1: Ages ranging 0-20, 2: Ages ranging 21-40, 3: Ages ranging 41-60, 4: Ages over 61.

