

The association between growth factors and blood factors with early childhood caries

Hoda Shamsaddin DDS, MSc¹, Fatemeh Jahanimoghadam DDS, MSc²,
Hamidreza Poureslami DDS, MSc³, Ali Akbar Haghdoost MD, PhD⁴

Original Article

Abstract

BACKGROUND AND AIM: Early childhood caries (ECC) is a specific form of rampant dental caries affecting infants and young children. ECC is still a problem that threatens the health of the community and its management is important. It is believed that the majorities of children with ECC also suffer from malnutrition, anemia, low weight and altered physical growth patterns. This study was undertaken to compare blood indices [mean corpuscular volume (MCV), hemoglobin and serum ferritin] in a population of children in Kerman, Iran, and the association between growth factors, blood parameters, and ECC.

METHODS: In this cross-sectional study, 240 children were selected from Afzalipour Hospital of Kerman. The subjects consisted of 2 to 6-year children, who needed blood sampling for different diagnostic reasons. Data were collected through clinical oral examinations, anthropometric measures, blood indices measurement (MCV, hemoglobin and serum ferritin) and structured questionnaire in order to assess demographic characteristics, nutritional habits and the presence or absence of systemic conditions. Data were analyzed using SPSS software.

RESULTS: The mean age of subjects was 50.79 months. Of 240 children included in this study, 124 (52.1%) were girls and the rest (47.9%) were boys. Statistical test revealed that there were statically significant differences in weight as well as the height of children in experimental and control groups ($P < 0.050$). No significant differences were detected in the frequencies of low hematocrit levels between the groups with and without dental caries ($P > 0.050$).

CONCLUSION: No significant association was observed between ECC and blood indices (MCV, hemoglobin and serum ferritin), but the mean height and weight in the caries-free group were significantly higher.

KEYWORDS: Dental Caries; Iron Deficiency; Anemia; Preschool Child; Growth Factors; Blood Factors; Early Childhood Caries

Citation: Shamsaddin H, Jahanimoghadam F, Poureslami H, Haghdoost AA. **The association between growth factors and blood factors with early childhood caries.** *J Oral Health Oral Epidemiol* 2017; 6(4): 196-202.

Dental caries is a transmissible infectious disease and its initiation and progression are affected by various factors.¹ It has been demonstrated that children with higher counts of *Streptococcus mutans* and *Lactobacilli* are 5 times more prone to have early childhood caries (ECC) as compared to those with lower

counts of these bacterial species.²

The American Academy of Pediatric Dentistry (AAPD) has defined ECC as the presence of one or several tooth surfaces (cavitated or non-cavitated) which have been lost or restored due to caries in any of the primary teeth in 71-month or younger children.³ It has been shown that children of

1- Assistant Professor, Oral and Dental Diseases Research Center AND Kerman Social Determinants on Oral Health Research Center AND Department of Pediatric Dentistry, School of Dentistry, Kerman University of Medical Sciences, Kerman, Iran

2- Associate Professor, Oral and Dental Diseases Research Center AND Kerman Social Determinants on Oral Health Research Center AND Department of Oral Medicine, School of Dentistry, Kerman University of Medical Sciences, Kerman, Iran

3- Professor, Oral and Dental Diseases Research Center AND Kerman Social Determinants on Oral Health Research Center AND Department of Oral Medicine, School of Dentistry, Kerman University of Medical Sciences, Kerman, Iran

4- Professor, Research Center for Modeling in Health, Institute for Futures Studies in Health, Kerman University of Medical Sciences, Kerman, Iran

Correspondence to: Fatemeh Jahanimoghadam DDS, MSc
Email: fatemehjehani4@gmail.com

families with low socioeconomic status, irrespective of culture and ethnicity, suffer from this condition more than other groups. In general and based on the studies carried out in almost all the countries, varying prevalence rates have been reported in different age groups.

ECC has been reported to be the most common chronic disease in children in the United States. Dental caries is five times more common than asthma and seven times more common than hay fever.⁴ Furthermore, the prevalence of ECC in Iran has been reported to be 3-44% in different parts of the country,⁵⁻⁷ which necessitates further research about the condition and evaluation of techniques to deal with it. The current knowledge on the effects of ECC on the nutritional status of children is not sufficient and it is believed that the majorities of children with ECC also suffer from malnutrition, anemia, low weight and altered physical growth patterns.⁸

Despite the epidemic nature of dental caries and iron deficiency, limited studies are available on the association between these two entities. ECC is still a problem that threatens the health of the community and its management is important. ECC might lead to the malocclusion of permanent teeth, phonetic problems, and low self-confidence. Continuous and unbearable pain in young children, such as a toothache, have serious and sometimes irreversible effects on the growth and development of children.⁹ Moreover, caries is more prevalent in deciduous teeth. This phenomenon itself is one of the most important factors in shifting caries to newly erupted permanent teeth.¹⁰ The children's health affects their nutrition, sleep, smile, speech, and socialization. In addition, the facial appearance affects the children's mental health and, subsequently the establishment of social relationships.¹¹ Furthermore, clinicians have also found that an attractive appearance and social acceptance can help the person in achieving social success.¹² Based on a research, weight gain and other parameters related to the

growth and development of children are normal when the teeth are healthy, which might be attributed to the efficacy of mastication and proper nutrition.¹³

ECC predisposes more severe dental conditions in future,¹⁴ and increases the sensitivity of children to somatic pains. Limitations in selecting food, decreased appetite, anorexia and a decrease in food enjoyment are complications that associate dental caries with weight loss, delayed growth and poor development.¹⁵⁻¹⁷ On the other hand, it has been demonstrated that the weight loss is compensated after the dental health has been improved.¹⁸ Therefore, identification of malnutrition risk factors results in a decrease in its occurrence, which in turn improves the children's health-related indices.

The present study was undertaken to evaluate growth and blood factors in children with ECC.¹⁹ Therefore, the possible role of some factors relevant to malnutrition in this type of dental caries would be elucidated.

Methods

This cross-sectional study was undertaken using clinical examination. A checklist, including demographic characteristics (age, sex) and the presence or absence of systemic conditions, was completed by an examiner (a trained pedodontist) on 240 children. All the 2 to 6-year children in the pediatric department of Afzalipour Hospital of Kerman, south-east of Iran, who needed blood sampling for different diagnostic reasons, were included in this study through random sampling. The parents were asked about children's feeding pattern during infancy and birth weight. The ethical code (K/88/07) was allocated to this study by the Oral Diseases Research Center of Kerman. The parents signed an informed consent form for the participation of their children in the study.

Children with ECC were compared with a control group of caries-free children. Two groups of children were included in this study. The inclusion criteria were as follows: no specific physical or systemic condition related

to ECC or iron deficiency as reported by the parents, absence of severe renal insufficiency and hypothyroidism (anemia is one of the complications of these two medical conditions), an age range of 2 to 6 years. Children with systemic disease, severe renal insufficiency and hypothyroidism were excluded.

The children underwent a clinical examination under a 100-W light using disposable dental mirrors and explorers and placed in one of the following groups. G1: caries-free, G2: decalcification on the maxillary incisors, G3: cavity on the maxillary incisors, and G4: decalcification on some maxillary incisors and cavities on some others.²⁰

The clinical evaluation of the nutritional status involves several tests to correctly determine the presence of malnutrition. The clinical parameters of children including height, weight and the upper mid-arm circumference (MAC) were measured. The MAC was measured on the right side. A tape measure, with no capacity to be stretched, was used for such measurement. To this end, the examiner measured the distance between the acromion process of the scapula and the elbow, determined the middle point, and then measured the circumference of the arm at this point without exerting any pressure on the arm.²¹ The most important blood tests that might indicate iron deficiency are hemoglobin, ferritin sodium and to a lesser extent mean corpuscular volume (MCV),²² which were

evaluated in the present study. Based on a definition by World Health Organization (WHO), a hemoglobin level under 11 or a hematocrit value under 33% in 6 to 59-month children, and hemoglobin levels under 11.5 or hematocrit levels under 34% in 5 to 11-year children indicate anemia.²³ If iron deficiency anemia is assessed based on serum ferritin levels, at ages under 5 and ages between 5 to 15 years, ferritin levels under 12 and 15 are considered abnormal, respectively.²⁴

Data were analyzed using SPSS software (version 20, IBM Corporation, Armonk, NY, USA). Mean \pm standard deviation (SD) and number (percentage) were used for data description. Analysis of variance (ANOVA), Student's independent t-test, and chi-square or Fisher's exact tests were used for data analysis. Statistical significance was set at $P < 0.050$.

Results

A total of 240 children were included in this cross-sectional study. 124 subjects (52.1%) were girls and the rest (47.9%) were boys. Table 1 shows frequency distribution of the children according to sex, feeding pattern, caries status, birth weight, and blood parameters. Demographic characteristics and selected blood tests between ECC and control groups have been shown in table 2. ANOVA test outcomes showed that only the mean height in the caries-free group was significantly higher than that in other groups (Table 3).

Table 1. Distribution of the children according to sex, feeding pattern, caries, birth weight, and blood parameters

| Parameter | | n (%) |
|----------------------|--|------------|
| Sex | Female | 124 (51.7) |
| | Male | 116 (48.3) |
| Feeding pattern | Breastfeeding | 166 (69.2) |
| | Bottle feeding | 36 (15.0) |
| | Both | 38 (15.8) |
| Blood test | Normal | 84 (35.0) |
| | Decreased | 156 (65.0) |
| Weight at birth time | Normal | 220 (91.7) |
| | Low birth weight | 20 (8.3) |
| Caries | Caries free | 83 (34.6) |
| | Decalcification of anterior upper teeth | 18 (7.5) |
| | Cavity on anterior upper teeth | 126 (52.5) |
| | Decalcification in some teeth and cavity in others | 13 (5.4) |

Table 2. Demographic characteristics and selected laboratory tests among 2 to 6-year Iranian children

| Characteristic | Caries | |
|------------------------------|---------------|---------------|
| | No (n = 83) | Yes (n = 159) |
| Ferritin (ng/ml) (mean ± SD) | 34.63 ± 19.16 | 34.39 ± 24.93 |
| Hb (g/dl) (mean ± SD) | 12.37 ± 1.02 | 12.50 ± 2.48 |
| Height (cm) (mean ± SD) | 99.53 ± 14.15 | 96.94 ± 9.97 |
| MAC (mean ± SD) | 15.09 ± 2.74 | 14.98 ± 1.66 |
| MCV (mean ± SD) | 77.61 ± 4.59 | 76.80 ± 5.64 |
| Weight (kg) (mean ± SD) | 16.34 ± 4.24 | 15.22 ± 3.29 |
| Age (month) (mean ± SD) | 51.87 ± 16.45 | 50.23 ± 14.42 |

SD: Standard deviation; Hb: Hemoglobin; MAC: Mid-arm circumference; MCV: Mean corpuscular volume

The 4 groups were converted to two groups, with and without caries, and re-analyzed. Student's independent t-test analysis showed that only the mean weight of children who had no caries was significantly higher than children with any type of caries.

In addition, Fisher's exact test did not exhibit any significant differences in the frequencies of low hematocrit values and low birth weight between the different age groups with caries. Similarly, chi-square test did not show any significant differences in the frequencies of low hematocrit values and low birth weight between the two groups (Table 4).

Based on the results, the mean height of caries-free children was significantly higher than the children with caries. The mean weight of caries-free children was significantly higher than that of children with caries. There were no significant differences in the frequencies of low hematocrit levels between groups with and without caries. The frequencies of low birth weight were not significantly different between groups with different caries rates. There were no

significant differences in the frequencies of low birth rates between groups with and without caries.

Discussion

At present, there are insufficient data on the relationship between caries and iron deficiency. Meanwhile, the available data are contradictory. In this study, children with ECC and caries-free children were included to evaluate the relationship between MCV, hemoglobin, and serum ferritin levels and dental caries. Height, weight and MAC of all participants were measured.

To date, only a limited number of studies have assessed the nutritional status of young children with ECC. In a study, the clinical criterion used for the diagnosis of ECC was the presence of smooth surface caries on 4 maxillary incisors.²⁵ In the present study, the clinical criterion for the diagnosis of ECC was the presence of smooth surface caries with cavities or decalcification on 4 maxillary incisors. Initial studies have shown that children with ECC might have lower mean weight, compared to caries-free children.

Table 3. The comparison of measured parameter between four groups according to caries stages

| Parameter | Caries free (n = 83) | Decalcification of anterior upper teeth (n = 18) | Cavity on anterior upper teeth (n = 126) | Decalcification in some teeth and cavity in others (n = 13) | P* |
|------------------------------|----------------------|--|--|---|-------|
| Weight (kg) (mean ± SD) | 16.34 ± 4.25 | 14.07 ± 2.33 | 15.41 ± 3.26 | 15.03 ± 4.47 | 0.060 |
| Height (cm) (mean ± SD) | 99.54 ± 14.15 | 93.28 ± 11.99 | 97.98 ± 9.43 | 91.65 ± 10.32 | 0.040 |
| MAC | 15.09 ± 2.74 | 14.36 ± 0.92 | 15.08 ± 1.66 | 15.00 ± 2.11 | 0.570 |
| Hb (g/dl) (mean ± SD) | 12.38 ± 1.02 | 12.52 ± 1.04 | 12.27 ± 1.12 | 12.46 ± 0.95 | 0.710 |
| Ferritin (ng/ml) (mean ± SD) | 34.63 ± 19.10 | 24.39 ± 13.93 | 35.62 ± 26.51 | 38.66 ± 18.78 | 0.240 |
| MCV | 77.61 ± 4.59 | 77.92 ± 6.28 | 76.58 ± 5.74 | 77.47 ± 3.87 | 0.480 |

*Analysis of variance (ANOVA)

SD: Standard deviation; MAC: Mid-arm circumference; Hb: Hemoglobin; MCV: Mean corpuscular volume

Table 4. The comparison of measured parameters between two groups [with and without ECC (early childhood caries)]

| Parameter | Caries | | P* |
|------------------------------|---------------|---------------|-------|
| | No (n = 83) | Yes (n = 157) | |
| Weight (kg) (mean ± SD) | 16.35 ± 4.25 | 15.23 ± 3.29 | 0.040 |
| Height (cm) (mean ± SD) | 99.54 ± 14.16 | 96.91 ± 9.99 | 0.130 |
| MAC | 15.09 ± 2.75 | 14.99 ± 1.64 | 0.740 |
| Hb (g/dl) (mean ± SD) | 12.38 ± 1.02 | 12.31 ± 1.10 | 0.660 |
| Ferritin (ng/ml) (mean ± SD) | 34.63 ± 19.16 | 34.58 ± 25.01 | 0.980 |
| MCV | 77.61 ± 4.59 | 76.81 ± 5.66 | 0.260 |

*Student's independent t-test

SD: Standard deviation; Hb: Hemoglobin; MAC: Mid-arm circumference; MCV: Mean corpuscular volume

Therefore, it has been suggested that these children might not have received adequate calories.^{26,27} In this study, only the weight of children, who were caries-free, was significantly higher than that in children with caries.

Thomas and Primosch compared the weight of children with ECC, with the reference norms for children, and concluded that children with ECC had no noticeable weight loss.²⁷ The limitation of that study and similar studies is the fact that they have used only the children's weight as a criterion for the evaluation of their nutritional status, which is not sufficient for the evaluation of the clinical nature of nutrition. On the other hand, Bagherian and Sadeghi evaluated the relationship between body mass index (BMI) and dental caries in preschool children, where they showed a higher rate of dental caries in overweight children. They attributed this problem to the possible relationship between a higher intake of high-calorie foodstuffs deficient in nutritive materials by these children.²⁸ Several studies support these findings.²⁹⁻³²

Similar to our results, Gaur and Nayak suggested that severe ECC negatively influenced the children weights. Awareness and parents' education may help in improving their weights.³³ Clarke et al.²² evaluated the nutritional status of children with ECC using different clinical criteria. The clinical measurements consisted of height, weight and MAC. In addition, they evaluated the blood samples of the subjects for hemoglobin, MCV and serum albumin and

ferritin levels. These measurements were compared with those in a control group. The results of the study suggested that ECC might be a risk maker for iron deficiency anemia. In their study, 80% of the subjects exhibited low serum levels of ferritin and in 28% of children, the hemoglobin levels were lower than the standard levels. Interestingly, after rendering dental treatments to children with ECC, their ferritin and hemoglobin levels were improved.²²

Shaoul et al. showed a significant relationship between ECC and low serum levels of ferritin.³⁴ However, the mechanism of the relationship between iron deficiency and dental caries is unknown. Several hypotheses have been proposed in this regard to date. A hypothesis, based on a study by Gaur and Nayak,³³ suggests that a decrease in hemoglobin levels can be associated with inflammatory responses in the human body. The inflammatory responses in ECC might result in the production of cytokines that inhibit the synthesis of erythrocytes and decrease hemoglobin levels. A decrease in hemoglobin levels is common in many chronic diseases. Furthermore, severe ECC (S-ECC) is known as a chronic disease.

Schroth et al. reported that children with ECC have altered dietary habits that can predispose them to nutritional deficiencies such as iron deficiency.³ Fretham hypothesized that iron deficiency in children leads to a decrease in mental capacities, a decrease in motor skills and an increase in anxiety levels. Early diagnosis of such

deficiency, for example by detecting ECC in children, might result in the early initiation of treatment.³⁵ Unlike these studies, our result did not depict any association between ECC and blood parameters.

One of the shortcomings of the present study was the inability to match the samples completely. Although a consistent age range was considered for all the samples and the subjects were evaluated separately in relation to their sex, socioeconomic matching was not possible. S-ECC is more prevalent in children with lower socioeconomic status³⁶ and it is more difficult to find caries-free subjects in these socioeconomic levels. According to the important role of various factors in the induction of S-ECC, such as family income and the parents' educational levels, it is not possible to extend the results of this study to the whole community. Despite the limitations of this study, our results added the association

investigation between ECC and blood parameters and developmental indices to the literature and provided additional evidence for the complexity of this relationship.

Conclusion

No significant associations between ECC and blood parameters were observed. However, lower height and weight were significant determinants of caries experience. Finally, it is suggested that height, weight, and diet of children with ECC should be more precisely evaluated periodically.

Conflict of Interests

Authors have no conflict of interest.

Acknowledgments

This study was supported by the Research Committee of Kerman University of Medical Sciences, Iran.

References

1. Casamassimo PS, Thikkurissy S, Edelstein BL, Maiorini E. Beyond the DMFT: The human and economic cost of early childhood caries. *J Am Dent Assoc* 2009; 140(6): 650-7.
2. Ramos-Gomez FJ, Weintraub JA, Gansky SA, Hoover CI, Featherstone JD. Bacterial, behavioral and environmental factors associated with early childhood caries. *J Clin Pediatr Dent* 2002; 26(2): 165-73.
3. Schroth RJ, Levi J, Kliewer E, Friel J, Moffatt ME. Association between iron status, iron deficiency anaemia, and severe early childhood caries: A case-control study. *BMC Pediatr* 2013; 13: 22.
4. Filstrup SL, Briskie D, da Fonseca M, Lawrence L, Wandera A, Inglehart MR. Early childhood caries and quality of life: Child and parent perspectives. *Pediatr Dent* 2003; 25(5): 431-40.
5. Askarizadeh N, Siyonat P. The prevalence and pattern of nursing caries in preschool children of Tehran. *J Indian Soc Pedod Prev Dent* 2004; 22(3): 92-5.
6. Bargrizan M, Rahimi M, Moghadam B. Nursing caries in 2-4 years old children in Tehran. An epidemiologic survey. *J Dent Sch Shahid Beheshti Univ Med Sci* 2001; 18(4): 9-15. [In Persian].
7. Poureslami H, Adhemi SH. Relationship between ECC and feeding habits among a group of babies & toddlers in Kerman. *Journal of Islamic Dental Association of Iran* 2001; 17: 47-55. [In Persian].
8. Schroth RJ, Jeal NS, Kliewer E, Sellers EA. The relationship between vitamin D and severe early childhood caries: A pilot study. *Int J Vitam Nutr Res* 2012; 82(1): 53-62.
9. Larson K, Russ SA, Crall JJ, Halfon N. Influence of multiple social risks on children's health. *Pediatrics* 2008; 121(2): 337-44.
10. Jahanimoghadam F, Poureslami H, Shamsaddin H, Horri A, Khazaeli P, Mahvi AH. Effect of ER: Yag laser on sodium fluoride varnish uptake by primary tooth enamel: An in-vitro study. *Research Report Fluoride* 2016; 49(4 Pt 2): 538-48.
11. Yusuf H, Gherunpong S, Sheiham A, Tsakos G. Validation of an English version of the Child-OIDP index, an oral health-related quality of life measure for children. *Health Qual Life Outcomes* 2006; 4: 38.
12. Jahanimoghadam F, Momenidanayee S, Karimifshar M. Correction of severe tooth rotation by using two different orthodontic appliances: Report of two cases. *J Oral Health Oral Epidemiol* 2016; 5(1): 46-51.
13. Monse B, Duijster D, Sheiham A, Grijalva-Eternod CS, van Palenstein Helderma W, Hobdell MH. The effects of extraction of pulpally involved primary teeth on weight, height and BMI in underweight Filipino children. A cluster randomized clinical trial. *BMC Public Health* 2012; 12: 725.
14. Greenwell AL, Johnsen D, DiSantis TA, Gerstenmaier J, Limbert N. Longitudinal evaluation of caries patterns from the primary to the mixed dentition. *Pediatr Dent* 1990; 12(5): 278-82.

15. Miller J, Vaughan-Williams E, Furlong R, Harrison L. Dental caries and children's weights. *J Epidemiol Community Health* 1982; 36(1): 49-52.
16. Ayhan H, Suskan E, Yildirim S. The effect of nursing or rampant caries on height, body weight and head circumference. *J Clin Pediatr Dent* 1996; 20(3): 209-12.
17. Benzian H, Monse B, Heinrich-Weltzien R, Hobdell M, Mulder J, van Palenstein, Helderma W. Untreated severe dental decay: A neglected determinant of low Body Mass Index in 12-year-old Filipino children. *BMC Public Health* 2011; 11: 558.
18. Ribeiro NM, Ribeiro MA. Breastfeeding and early childhood caries: A critical review. *J Pediatr (Rio J)* 2004; 80(5 Suppl): S199-S210.
19. Sullivan DH. The role of nutrition in increased morbidity and mortality. *Clin Geriatr Med* 1995; 11(4): 661-74.
20. Shivakumar K, Prasad S, Chandu G. International caries detection and assessment system: A new paradigm in detection of dental caries. *J Conserv Dent* 2009; 12(1): 10-6.
21. Committee on Nutrition AAO. Appendix I. Procedures for measuring growth parameters. Elk Grove Village, IL: American Academy of Pediatrics; 1998. p. 168-74
22. Clarke M, Locker D, Berall G, Pencharz P, Kenny DJ, Judd P. Malnourishment in a population of young children with severe early childhood caries. *Pediatr Dent* 2006; 28(3): 254-9.
23. Scott SP, Chen-Edinboro LP, Caulfield LE, Murray-Kolb LE. The impact of anemia on child mortality: An updated review. *Nutrients* 2014; 6(12): 5915-32.
24. Jose B, King NM. Early childhood caries lesions in preschool children in Kerala, India. *Pediatr Dent* 2003; 25(6): 594-600.
25. Acs G, Lodolini G, Kaminsky S, Cisneros GJ. Effect of nursing caries on body weight in a pediatric population. *Pediatr Dent* 1992; 14(5): 302-5.
26. Miller J, Vaughan-Williams E, Furlong R, Harrison L. Dental caries and children's weights. *J Epidemiol Community Health* 1982; 36(1): 49-52.
27. Thomas CW, Primosch RE. Changes in incremental weight and well-being of children with rampant caries following complete dental rehabilitation. *Pediatr Dent* 2002; 24(2): 109-13.
28. Bagherian A, Sadeghi M. Association between dental caries and age-specific body mass index in preschool children of an Iranian population. *Indian J Dent Res* 2013; 24(1): 66-70.
29. Norberg C, Hallstrom SU, Matsson L, Thorngren-Jerneck K, Klingberg G. Body mass index (BMI) and dental caries in 5-year-old children from southern Sweden. *Community Dent Oral Epidemiol* 2012; 40(4): 315-22.
30. Trikaliotis A, Boka V, Kotsanos N, Karagiannis V, Hassapidou M. Short communication: Dmfs and BMI in preschool Greek children. An epidemiological study. *Eur Arch Paediatr Dent* 2011; 12(3): 176-8.
31. Vazquez-Nava F, Vazquez-Rodriguez EM, Saldivar-Gonzalez AH, Lin-Ochoa D, Martinez-Perales GM, Joffre-Velazquez VM. Association between obesity and dental caries in a group of preschool children in Mexico. *J Public Health Dent* 2010; 70(2): 124-30.
32. Willershausen B, Moschos D, Azrak B, Blettner M. Correlation between oral health and body mass index (BMI) in 2071 primary school pupils. *Eur J Med Res* 2007; 12(7): 295-9.
33. Gaur S, Nayak R. Underweight in low socioeconomic status preschool children with severe early childhood caries. *J Indian Soc Pedod Prev Dent* 2011; 29(4): 305-9.
34. Shaoul R, Gaitini L, Kharouba J, Darawshi G, Maor I, Somri M. The association of childhood iron deficiency anaemia with severe dental caries. *Acta Paediatr* 2012; 101(2): e76-e79.
35. Fretham SJ, Carlson ES, Georgieff MK. The role of iron in learning and memory. *Adv Nutr* 2011; 2(2): 112-21.
36. Costa LR, Daher A, Queiroz MG. Early childhood caries and body mass index in young children from low income families. *Int J Environ Res Public Health* 2013; 10(3): 867-78.